

## A NOVEL NRF2-MEDIATED ANTIOXIDATIVE AND ANTI-INFLAMMATORY MECHANISM THROUGH FERROPORTIN 1 INDUCTION: A POTENTIAL CYTOPROTECTIVE MECHANISM AGAINST ISCHEMIA-REPERFUSION INJURIES

Nobuhiko Harada<sup>1)</sup>, Shin Matsuura<sup>1)</sup>, Masaya Kanayama<sup>2)</sup>, Aruto Yoshida<sup>2)</sup> and Ken Itoh<sup>1)</sup>

**Abstract** The long-term function of a graft after transplantation is dependent upon ischemia-reperfusion (I/R) injury, which represents the most important alloantigen-independent damage processes during transplantation. Basic leucine zipper (b-Zip) transcription factor NF-E2-related factor 2 (Nrf2) is protective against I/R injury by promoting antioxidative and anti-inflammatory mechanisms via up-regulation of antioxidant-responsive element (ARE)-regulated genes such as heme oxygenase 1 and  $\gamma$ -glutamylcysteine ligase. We previously demonstrated that macrophage Nrf2 plays an important role in an elastase-induced acute lung injury model. To further clarify the mechanisms that act downstream of Nrf2, we performed microarray analysis in the RAW264.7 murine macrophage cell line using endogenous electrophilic prostaglandin 15-deoxy- $\Delta^{12,14}$ -prostaglandin J<sub>2</sub> (15d-PGJ<sub>2</sub>). As a result, we identified ferroportin 1 (Fpn1) as a novel Nrf2 target gene. Fpn1 is the sole iron exporter in mammals and regulates iron homeostasis. On the other hand, free intracellular iron is detrimental to cells by enhancing both inflammation and Fenton reaction-mediated oxidative stress. We demonstrated that Nrf2 activation by several electrophilic compounds commonly result in the up-regulation of *Fpn1* mRNA in bone marrow-derived and peritoneal macrophages obtained from wild-type, but not from *Nrf2* knockout mice. Furthermore, Nrf2 activation enhances iron release from the murine macrophage cell line J774.1. It is previously reported that LPS suppresses *Fpn1* mRNA expression, which leads to iron retention in monocytes and macrophages. We showed that while LPS suppresses *Fpn1* mRNA expression in human macrophages, Nrf2 activation restores the expression of *Fpn1*. Thus, we propose that iron-metabolism regulation may be an important factor of the Nrf2-mediated cytoprotective mechanism.

Hirosaki Med. J. 64, Supplement : S74—S80, 2013

**Key words:** Inflammation; Oxidative stress; Iron; Nrf2

### 1. Iron homeostasis is important for normal cell function.

Iron is an essential micronutrient for cell survival and proliferation in mammal. Excess cellular free iron generates reactive oxygen species (ROS) through the Fenton reaction and damage lipid membranes, proteins, and nucleic acids, while iron deficiency affects cell growth and leads to cell death<sup>1-3)</sup>. Therefore, the cellular iron level is strictly regulated at the levels of

uptake, storage and export.

### 2. Ferroportin 1 is the sole iron exporter in mammalian cells.

Ferroportin 1 (Fpn1) is the sole non-heme iron exporter in mammals and is highly expressed in duodenal enterocytes and tissue macrophages<sup>1)</sup>. Fpn1 is localized to the basolateral membranes of duodenal enterocytes and plays an important role in iron absorption from diet<sup>4)</sup>. Since dietary iron

<sup>1)</sup>Department of Stress Response Science, Hirosaki University Graduate School of Medicine

<sup>2)</sup>Central Laboratories for Frontier Technology, Kirin Holdings Co. Ltd.

Corresponding author; Ken Itoh, MD & PhD

Department of Stress Response Science, Hirosaki

University Graduate School of Medicine, 5 Zaifucho, Hirosaki 036-8562, Japan

Phone: +81-172-39-5158

Fax: +81-172-39-5157

E-mail: itohk@cc.hirosaki-u.ac.jp

intake is restricted, a large amount of iron for daily requirement is recycled from the senescent red blood cells by the reticuloendothelial macrophages<sup>1, 2</sup>. After the phagocytosis of senescent erythrocytes by macrophages, the heme-derived iron is subsequently exported to circulation by the cooperation of heme oxygenase 1 (HO-1) and Fpn1<sup>4</sup>. Recently, it is suggested that Fpn1 is also expressed in the brain and might be involve in iron export from neural cells<sup>5, 6</sup>. Fpn1 expression is controlled by both transcriptionally and post-transcriptionally. *Fpn1* mRNA expression levels are transcriptionally regulated by erythrophagocytosis or iron overload in macrophages<sup>7-9</sup>, but the mechanisms are not completely understood. Post-transcriptionally, Fpn1 protein expression is regulated either through translational regulation by an iron-responsive element (IRE) within the mRNA<sup>10</sup> or through protein degradation by hepcidin, a peptide hormone for iron homeostasis mainly produced by hepatocytes in response to iron, inflammatory stimuli, and hypoxia<sup>11-13</sup>.

### **3. Transcription factor Nrf2 regulates cytoprotective genes.**

NF-E2-related factor 2 (Nrf2) is a basic leucine zipper (b-Zip) transcription factor and plays an important role in the cytoprotection against oxidative stress through the induction of antioxidant responsive element (ARE) -regulated genes<sup>14</sup>. Under unstimulated conditions, Nrf2 is bound by Kelch-like ECH-associated protein 1 (Keap1) in the cytoplasm and subject to the proteasomal degradation. The cysteine residues in Keap1 are modified by oxidative stress or electrophiles and Nrf2 is liberated from Keap1-mediated degradation. Subsequently, Nrf2 translocates from the cytoplasm into the nucleus and binds to AREs in the regulatory regions of the genes that encode antioxidant enzymes that protect the cell from the oxidative

damage. Thus, Nrf2 protein level is regulated by Keap1-mediated degradation process, and the stabilization of Nrf2 is required for the cellular protective response to oxidative and electrophilic stresses<sup>15, 16</sup>. Many bioactive food components such as sulforaphane (SFN), an isothiocyanate from broccoli sprouts, resveratrol from grapes, curcumin from turmeric, and carnosic acid from rosemary also activate Nrf2<sup>17-20</sup>. Nrf2 is ubiquitously expressed in mammalian tissues and Nrf2/ARE system regulates a battery of cytoprotective genes such as drug-detoxifying enzyme (glutathione S-transferase (GST) and NAD(P)H:quinone oxidoreductase 1 (NQO1)), antioxidant defense enzymes such as HO-1, enzymes used in glutathione synthesis (subunits of glutamate-cysteine ligase), enzymes for NADPH generation (malic enzyme, etc), and phase III drug transporters<sup>21</sup> (Figure 1). Ablation of Nrf2 increases the cellular sensitivity to oxidative stress and inflammation<sup>22-24</sup>. The activation of Nrf2 signaling pathway has a protective effect against oxidative stress including ischemia-reperfusion injury and inflammation in the brain<sup>25, 26</sup>.

### **4. Activation of Nrf2 up-regulates *Fpn1* mRNA expression and enhances iron efflux through Fpn1 induction.**

Recently, we and others independently reported that Nrf2 regulates *Fpn1* mRNA expression in macrophages<sup>27, 28</sup>. One report comes from the study regarding the heme-mediated regulation of iron homeostasis. It was previously reported that heme activates *Fpn1* transcription in an iron-independent manner during erythrophagocytosis<sup>29</sup>. Marro S. *et al.* found that heme-mediated *Fpn1* mRNA induction is achieved through Nrf2 and BTB and CNC Homology 1 (Bach1)<sup>27</sup>. Bach1 also belongs to the CNC transcription factor family. It bind to ARE-like sequences and acts as a




Figure 1 Nrf2/ARE system regulates a battery of cytoprotective genes.

transcriptional repressor in a battery of ARE-regulated genes including *HO-1*<sup>30)</sup>. Bach1 export from the nucleus and its protein degradation are enhanced by heme<sup>31)</sup>. Thus, heme inhibits the repressor function of Bach1 and enhances *HO-1* gene expression. Furthermore, Nrf2 is also activated by heme and enhances *HO-1* gene expression. According to Marro *et al.*, heme also increases *Fpn1* mRNA in the RAW264.7 murine macrophage cell line and heme-mediated *Fpn1* up-regulation is controlled by an ARE element approximately 7kb upstream from the *Fpn1* transcription start site<sup>28)</sup>. They also showed that Nrf2 positively regulates *Fpn1* promoter activity while Bach1 negatively regulates its activity. Thus, they hypothesize that iron recycling in macrophages are transcriptionally co-regulated by heme through Bach1 and Nrf2. We have been interested in Nrf2-mediated anti-inflammatory mechanisms. We previously demonstrated that macrophage Nrf2 plays a critical role in an elastase-induced acute lung injury model<sup>32)</sup>. To further elucidate the

mechanisms acting downstream of Nrf2, we performed microarray analysis in RAW264.7 macrophages using endogenous electrophilic prostaglandin 15d-PGJ<sub>2</sub>. As a result, we identified *Fpn1* as a novel Nrf2 target gene<sup>27)</sup>. Nrf2 activation by several electrophilic compounds commonly resulted in the up-regulation of *Fpn1* mRNA in macrophages derived from wild-type mice but not from *Nrf2* knockout mice. Further, Nrf2 activation enhanced iron release from the murine macrophage cell line J774.1. Previous investigations showed that inflammatory stimuli, such as LPS, down-regulates macrophage *Fpn1* expression by transcriptional and hepcidin-mediated post-translational mechanisms leading to iron sequestration by macrophages<sup>33-36)</sup>. We showed that two Nrf2 activators, diethyl maleate and SFN, restored the LPS-induced suppression of *Fpn1* mRNA in macrophages<sup>27)</sup>. These results suggest that Nrf2 regulates iron efflux from cells through *Fpn1* gene transcription and Nrf2 may control iron metabolism during inflammation. Recent studies revealed that the

decrease of intramacrophage iron levels due to Fpn1 up-regulation attenuates TLR4-mediated inflammation by inhibiting expression of pro-inflammatory cytokines, such as IL-6, TNF $\alpha$ , and INF- $\beta$ <sup>37, 38</sup>.

### 5. An emerging role of Fpn1 as an anti-inflammatory factor

In addition to the above-mentioned anti-inflammatory role of Fpn1 through iron metabolism, it is recently reported that Fpn1 activates intracellular signaling for inflammatory gene expression upon hepcidin binding<sup>39</sup>. Hepcidin induces Fpn1 degradation and internalization *via* activation of Jak2 and the subsequent phosphorylation of Fpn1. De Domenico *et al.* demonstrated that Jak2 activated upon hepcidin binding also phosphorylates Stat 3 that results in the upregulation of genes such as *Interleukin (IL) 7*, *IL17 receptor* and *SOCS*. Hepcidin pretreatment inhibits LPS-induced TNF $\alpha$  and IL-6 gene expression via SOCS induction. *In vivo*, hepcidin pretreatment rescues LPS-induced lethality.


### 6. Nrf2 may have a beneficial effect for ischemia-reperfusion injury through Fpn1 induction.

Ischemia-reperfusion injury occurs when blood flow to a tissue is disturbed and subsequently recanalized. It is related to a various disease states including myocardial infarction, stroke, and post-transplantation injury<sup>40-42</sup>. Hypoxia during ischemia and subsequent reoxygenation by reperfusion are thought to be the major factors contributing to ROS production and the subsequent cellular damage<sup>43</sup>. Iron is believed to play a critical role in neuronal injuries caused by oxidative stress in cerebral ischemia<sup>44</sup>. In the brain, it is well known that free iron significantly increases during ischemia and is responsible for oxidative damage. Increased amount of free

iron and ferritin have been observed in ischemic brain<sup>45, 46</sup>, while the iron chelator desferrioxamine (DFO) diminish the damaging role of free iron in the ischemic brain<sup>47, 48</sup>. Recently, Ding *et al.* have shown that ischemia increases intracellular iron contents by reducing Fpn1 expression in the brain<sup>49</sup>. Furthermore, Leonard *et al.* demonstrated that reoxygenation-specific activation of the Nrf2 mediates cytoprotective gene expression in ischemia-reperfusion injury<sup>50</sup>. Thus, we propose that the regulation of iron metabolism may be an important aspect of the Nrf2-mediated cytoprotective mechanism and it may have the protective role against ischemia-reperfusion injury (Figure 2).

### References

- 1) Hentze MW, Muckenthaler MU, Andrews NC. Balancing acts: molecular control of mammalian iron metabolism. *Cell* 2004;117:285-97.
- 2) Muñoz M, Villar I, García-Erce JA. An update on iron physiology. *World J Gastroenterol* 2009;15: 4617-26.
- 3) Yu Y, Kovacevic Z, Richardson DR. Tuning cell cycle regulation with an iron key. *Cell Cycle* 2007;6:1982-94.
- 4) Donovan A, Lima CA, Pinkus JL, et al. The iron exporter ferroportin/Slc40a1 is essential for iron homeostasis *Cell Metab* 2005;1:191-200.
- 5) Chen Y, Qian ZM, Du J, et al. Iron loading inhibits ferroportin1 expression in PC12 cells. *Neurochem Int* 2005;47:507-13.
- 6) Moos T, Rosengren Nielsen T, Skjørringe T, Morgan EH. Iron trafficking inside the brain. *J Neurochem* 2007;103:1730-40.
- 7) Delaby C, Pilard N, Hetet G, et al. A physiological model to study iron recycling in macrophages. *Exp Cell Res* 2005;310:43-53.
- 8) Knutson MD, Vafa MR, Haile DJ, Wessling-Resnick M. Iron loading and erythrophagocytosis increase ferroportin 1 (FPN1) expression in J774 macrophages. *Blood* 2003;102:4191-7.



**Figure 2** Nrf2 might be capable of improving iron homeostasis during I/R-induced inflammation by increasing *Fpn1* transcription along with the genes that have anti-inflammatory functions.

- 9) Delaby C, Pilard N, Puy H, Canonne-Hergaux F. Sequential regulation of ferroportin expression after erythrophagocytosis in murine macrophages: early mRNA induction by haem, followed by iron-dependent protein expression. *Biochem J* 2008;411:123-31.
- 10) Lymboussaki A, Pignatti E, Montosi G, Garuti C, Haile DJ, Pietrangelo A. The role of the iron responsive element in the control of ferroportin1/IREG1/MTP1 gene expression. *J Hepatol* 2003;39:710-5.
- 11) Nemeth E, Tuttle MS, Powelson J, et al. Hepcidin regulates cellular iron efflux by binding to ferroportin and inducing its internalization. *Science* 2004;306:2090-3.
- 12) Ganz T, Nemeth E. Regulation of iron acquisition and iron distribution in mammals. *Biochim Biophys Acta* 2006;1763:690-9.
- 13) Nicolas G, Chauvet C, Viatte L, et al. The gene encoding the iron regulatory peptide hepcidin is regulated by anemia, hypoxia, and inflammation. *J Clin Invest* 2002;110:1037-44.
- 14) Itoh K, Chiba T, Takahashi S, et al. An Nrf2/small Maf heterodimer mediates the induction of phase II detoxifying enzyme genes through antioxidant response elements. *Biochem Biophys Res Commun* 1997;236:313-22.
- 15) Kensler TW, Wakabayashi N, Biswal S. Cell survival responses to environmental stresses via the Keap1-Nrf2-ARE pathway. *Annu Rev Pharmacol Toxicol* 2007;47:89-116.
- 16) Taguchi K, Motohashi H, Yamamoto M. Molecular mechanisms of the Keap1-Nrf2 pathway in stress response and cancer evolution. *Genes Cells* 2011;16:123-40.
- 17) Fahey JW, Zhang Y, Talalay P. Broccoli sprouts: an exceptionally rich source of inducers of enzymes that protect against chemical carcinogens. *Proc Natl Acad Sci USA* 1997;94:10367-72.
- 18) Rubiolo JA, Mithieux G, Vega FV. Resveratrol protects primary rat hepatocytes against oxidative stress damage: activation of the Nrf2 transcription factor and augmented activities of antioxidant enzymes. *Eur J Pharmacol* 2008;591:66-72.
- 19) Balogun E, Hoque M, Gong P, et al. Curcumin activates the haem oxygenase-1 gene via regulation of Nrf2 and the antioxidant-responsive

- element. *Biochem J* 2003;371:887-95.
- 20) Kosaka K, Mimura J, Itoh K, et al. Role of Nrf2 and p62/ZIP in the neurite outgrowth by carnosic acid in PC12h cells. *J Biochem.* 2010;147:73-81.
  - 21) Kobayashi M, Yamamoto M. Nrf2-Keap1 regulation of cellular defense mechanisms against electrophiles and reactive oxygen species. *Adv Enzyme Regul* 2006;46:113-40.
  - 22) Thimmulappa RK, Lee H, Rangasamy T, et al. Nrf2 is a critical regulator of the innate immune response and survival during experimental sepsis. *J Clin Invest* 2006;116:984-95.
  - 23) Innamorato NG, Rojo AI, García-Yagüe AJ, Yamamoto M, de Ceballos ML, Cuadrado A. The transcription factor Nrf2 is a therapeutic target against brain inflammation. *J Immunol* 2008; 181:680-9.
  - 24) Cho HY, Jedlicka AE, Reddy SP, et al. Role of NRF2 in protection against hyperoxic lung injury in mice. *Am J Respir Cell Mol Biol* 2002;26:175-82.
  - 25) Satoh T, Kosaka K, Itoh K, et al. Carnosic acid, a catechol-type electrophilic compound, protects neurons both in vitro and in vivo through activation of the Keap1/Nrf2 pathway via S-alkylation of targeted cysteines on Keap1. *J Neurochem* 2008;104:1116-31.
  - 26) Zhao X, Sun G, Zhang J, et al. Transcription factor Nrf2 protects the brain from damage produced by intracerebral hemorrhage. *Stroke* 2007;38:3280-6.
  - 27) Harada N, Kanayama M, Maruyama A, et al. Nrf2 regulates ferroportin 1-mediated iron efflux and counteracts lipopolysaccharide-induced ferroportin 1 mRNA suppression in macrophages. *Arch Biochem Biophys* 2011;508:101-9.
  - 28) Marro S, Chiabrando D, Messana E, et al. Heme controls ferroportin1 (FPN1) transcription involving Bach1, Nrf2 and a MARE/ARE sequence motif at position -7007 of the FPN1 promoter. *Haematologica* 2010;95:1261-8.
  - 29) Delaby C, Pilard N, Puy H, Canonne-Hergaux F. Sequential regulation of ferroportin expression after erythrophagocytosis in murine macrophages: early mRNA induction by haem, followed by iron-dependent protein expression. *Biochem J* 2008;411:123-31.
  - 30) Jyrkkänen HK, Kuosmanen S, Heinäniemi M, et al. Novel insights into the regulation of antioxidant-response-element-mediated gene expression by electrophiles: induction of the transcriptional repressor BACH1 by Nrf2. *Biochem J* 2011;440: 167-74.
  - 31) Zenke-Kawasaki Y, Dohi Y, Katoh Y, et al. Heme induces ubiquitination and degradation of the transcription factor Bach1. *Mol Cell Biol* 2007;27:6962-71.
  - 32) Ishii Y, Itoh K, Morishima Y, et al. Transcription factor Nrf2 plays a pivotal role in protection against elastase-induced pulmonary inflammation and emphysema. *J Immunol* 2005;175:6968-75.
  - 33) Nemeth E, Rivera S, Gabayan V, et al. IL-6 mediates hypoferrremia of inflammation by inducing the synthesis of the iron regulatory hormone hepcidin. *J Clin Invest* 2004;113:1271-6.
  - 34) Weiss G, Goodnough LT. Anemia of chronic disease. *N Engl J Med* 2005;352:1011-23.
  - 35) Ludwiczek S, Aigner E, Theurl I, Weiss G. Cytokine-mediated regulation of iron transport in human monocytic cells. *Blood* 2003;101:4148-54.
  - 36) Liu XB, Nguyen NB, Marquess KD, Yang F, Haile DJ. Regulation of hepcidin and ferroportin expression by lipopolysaccharide in splenic macrophages. *Blood Cells Mol Dis* 2005;35:47-56.
  - 37) Wang L, Harrington L, Trebicka E, et al. Selective modulation of TLR4-activated inflammatory responses by altered iron homeostasis in mice. *J Clin Invest* 2009;119:3322-8.
  - 38) Wang L, Johnson EE, Shi HN, Walker WA, Wessling-Resnick M, Cherayil BJ. Attenuated inflammatory responses in hemochromatosis reveal a role for iron in the regulation of macrophage cytokine translation. *J Immunol* 2008;181:2723-31.
  - 39) De Domenico I, Zhang TY, Koenig CL, et al. Hepcidin mediates transcriptional changes that modulate acute cytokine-induced inflammatory responses in mice. *J Clin Invest* 2010;120:2395-405.
  - 40) Moens AL, Claeys J, Timmermans JP, Vrints CJ.

- Myocardial ischemia/reperfusion-injury, a clinical view on a complex pathophysiological process. *Int. J. Cardiol* 2005;100:179-90.
- 41) Zheng Z, Lee JE, Yenari MA. Stroke: molecular mechanisms and potential targets for treatment. *Curr. Mol. Med* 2003;3:361-72.
- 42) Husted TL, Lentsch AB. Anti-inflammatory approaches to the prevention of ischemia/reperfusion injury in solid organ transplantation. *Curr. Opin. Invest. Drugs*. 2005;6:508-12.
- 43) Li C, Jackson RM. Reactive species mechanisms of cellular hypoxia-reoxygenation injury. *Am. J. Physiol* 2002;82:C227-C241
- 44) Selim MH, Ratan RR. The role of iron neurotoxicity in ischemic stroke. *Ageing Res Rev* 2004;3:345-53.
- 45) Lipscomb DC, Gorman LG, Traystman RJ, Hurn PD. Low molecular weight iron in cerebral ischemic acidosis in vivo. *Stroke* 1998;29:487-92.
- 46) Chi SI, Wang CK, Chen JJ, Chau LY, Lin TN. Differential regulation of H- and L-ferritin messenger RNA subunits, ferritin protein and iron following focal cerebral ischemia-reperfusion. *Neuroscience* 2000;100:475-84.
- 47) Mu D, Chang YS, Vexler ZS, Ferriero DM. Hypoxia-inducible factor 1alpha and erythropoietin upregulation with deferoxamine salvage after neonatal stroke. *Exp Neurol* 2005;195:407-15.
- 48) Hanson LR, Roeytenberg A, Martinez PM, et al. Intranasal deferoxamine provides increased brain exposure and significant protection in rat ischemic stroke. *J Pharmacol Exp Ther* 2009; 330:679-86.
- 49) Ding H, Yan CZ, Shi H, et al. Hepcidin is involved in iron regulation in the ischemic brain. *PLoS One* 2011;6:e25324.
- 50) Leonard MO, Kieran NE, Howell K, et al. Reoxygenation-specific activation of the antioxidant transcription factor Nrf2 mediates cytoprotective gene expression in ischemia-reperfusion injury. *FASEB J* 2006;20:2624-6.