

Anatomical classification of breast sentinel lymph nodes using computed tomography-

lymphography

(CT リンフォグラフィーによる乳腺センチネルリンパ節の解剖学的分類)

申 請 者      弘前大学大学院医学研究科

総合医療・健康科学領域 放射線診断学教育研究分野

氏      名      藤田 環

指導教授      青木 昌彦

**Mini-abstract**

Information of the sentinel lymph nodes in patients with breast cancer that show wide variation in the number, branching patterns, location and flow direction can be obtained by Computed tomography-lymphography.

**Anatomical classification of breast sentinel lymph nodes using computed tomography-  
lymphography**

**Tamaki Fujita<sup>1</sup>, Hiroyuki Miura<sup>1</sup>, Hiroko Seino<sup>1,2</sup>, Shuichi Ono<sup>1</sup>, Takashi Nishi<sup>3</sup>, Akimasa  
Nishimura<sup>3</sup>, Kenichi Hakamada<sup>3</sup>, Masahiko Aoki<sup>1</sup>**

<sup>1</sup> Department of Radiology and Radiation Oncology, Graduate School of Medicine, Hirosaki  
University, 5 Zaifu-cho, Hirosaki-shi, Aomori 036-8562, Japan

<sup>2</sup> Department of Pathology and Bioscience, Graduate School of Medicine, Hirosaki University, 5  
Zaifu-cho, Hirosaki-shi, Aomori 036-8562, Japan

<sup>3</sup> Department of Gastroenterological Surgery, Graduate School of Medicine, Hirosaki University, 5  
Zaifu-cho, Hirosaki-shi, Aomori 036-8562, Japan

Correspondence to: Dr. Hiroyuki Miura, Department of Radiology and Radiation Oncology, Graduate  
School of Medicine, Hirosaki University, 5 Zaifu-cho, Hirosaki-shi, Aomori 036-8562, Japan

Phone number: +81-172-39-5103

Fax number: +81-172-33-5627

Email: [miuradio@hirosaki-u.ac.jp](mailto:miuradio@hirosaki-u.ac.jp)

**short running title** Anatomy of breast sentinel lymph by CT

**Original article**

## **Abstract**

*Purpose* To evaluate the anatomical classification and location of breast sentinel lymph nodes using computed tomography-lymphography.

*Materials and methods* Preoperative computed tomography-lymphography examinations for sentinel lymph nodes in 464 cases that were clinically diagnosed with node-negative breast cancer between July 2007 and June 2016 were retrospectively reviewed. Anatomical classification was performed based on the numbers of lymphatic routes and sentinel lymph nodes, the flow direction of lymphatic routes, and sentinel lymph nodes location.

*Results* Of the 464 cases reviewed, anatomical classification could be performed in 434 (93.5%), the largest number of cases showed single route/single sentinel lymph node (n = 296, 68.2%), followed by multiple routes/multiple sentinel lymph nodes (n = 59, 13.6%), single route/multiple sentinel lymph nodes (n = 53, 12.2%), and multiple routes/single sentinel lymph node (n = 26, 6.0%). A classification based on the flow direction of lymphatic routes showed that 429 cases (98.8%) had outward flow on the superficial fascia toward axillary lymph nodes, whereas a classification based on the height of sentinel lymph nodes showed that 323 cases (74.4%) belonged to the upper pectoral group of the axillary lymph nodes.

*Conclusion* There was wide variation in the number of lymphatic routes and their branching patterns and in the number, location, and direction of flow of sentinel lymph nodes. It is clinically very important to preoperatively understand the anatomical morphology of lymphatic routes and sentinel lymph nodes for optimal treatment of breast cancers, and computed tomography-lymphography is suitable for this purpose.

**Keywords** Anatomical classification, Axillary lymph node, Breast cancer, Computed tomography

lymphography, Sentinel lymph node

## Introduction

The sentinel lymph node (SLN) is any lymph node that directly receives lymph drainage from a tumor site (Uren et al. 2003). Uren et al. (2003) reported that SLNs need not necessarily be the node closest to the primary site and that lymphatic routes can bypass many other nodes before reaching the SLN.

Conventional methods for the detection of SLNs include the dye method and the radioisotope method. The dye method requires a high level of technical skills to trace the dye-stained lymphatic route to SLNs and can only be performed with support from a nuclear medicine department, while the scintigrams in the radioisotope method cannot clearly visualize the direct connection between primary SLNs and their afferent lymphatic routes (Yamamoto et al. 2016).

Computed tomography lymphography (CT-LG) involves CT after locally injecting a contrast agent. It can be conveniently performed, and the morphology and location of both lymphatic routes and SLNs can be visualized in detail, with minimal invasiveness (Yamamoto et al. 2016).

Yamamoto et al. (Yamamoto et al. 2016) used CT-LG to classify 549 cases into four anatomical categories, based on the numbers of lymphatic routes and lymph nodes: (a) single route/single SLN (355 cases, 65%; single lymphatic route and lymph node); (b) single route/multiple SLNs (62 cases, 11%; single lymphatic route with multiple lymph nodes); (c) multiple routes/single SLN (59 cases, 11%; multiple lymphatic routes with one lymph node); and (d) multiple routes/multiple SLNs (73 cases, 13%; multiple lymphatic routes and lymph nodes).

SLN biopsy is the gold standard procedure for axillary lymph node evaluation in breast cancer, and axillary lymph node dissection (ALND) is usually avoided in SLN-negative patients (Krag et al. 2010; The Japanese breast cancer society 2016; Lyman et al. 2005; Lyman et al. 2014). Thus, it is important to completely understand the anatomy and number of lymphatic routes and SLNs and their location. Location and flow direction of SLNs have been studied in cadavers (Suami et al. 2009a, Suami et al. 2009b) and radioisotope method (Blumgart et al. 2011a, Blumgart et al. 2011b), but anatomical morphology of lymphatic routes and SLNs using CT-LG has not been extensively studied since the report of Yamamoto et al. (Yamamoto et al. 2016). Thus, we decided to investigate them with CT-LG.



## **Materials and methods**

### **Patients**

This retrospective study was approved by our ethics committee, and informed consent for using case information was obtained from all patients. Their anonymity is preserved.

Preoperative examinations for SLNs in 480 cases with clinically node-negative breast cancer diagnosed between July 2007 and June 2016 at our institution were retrospectively reviewed. Six cases with a history of surgery and 10 with a history of chemotherapy were excluded. The remaining 464 cases (100%) were women, and the patients' characteristics are shown in Table 1.

### **CT-LG procedure**

CT-LG was performed using a 4-detector row CT scanner (Light Speed QX/i, GE Healthcare, Milwaukee, WI) until February 7, 2011, using a 64-detector row CT scanner (Somatom Definition, Siemens Healthcare, Munich, Germany) from February 8 to 13, 2011, and using a 64-detector row CT scanner (Discovery CT750 HD, GE Healthcare, Milwaukee, WI) from February 14, 2011 onward, with the following parameters: 120 kV; 200-400 mA; scan time, 0.5 s; and slice thickness, 1.25 mm (Light Speed QX/i), 1.00 mm (Somatom Definition), and 0.63 mm (Discovery CT750 HD). A mixture of 0.5 ml of 1%

lidocaine and 1 ml of iohexol (Omnipaque 300; Daiichi-Sankyo Company, Tokyo, Japan) or a mixture of 1.5 ml of 1% lidocaine and 1.5 ml of iohexol was injected into the nipple followed by gentle massage for 10 seconds. Each patient was placed in the supine position, with arms positioned in the cranial direction, as during breast cancer surgery. Scanning was initiated 60 seconds after contrast agent injection. Scans were appropriately added after 3 minutes, depending on the contrast spread.

All CT-LG data sets were loaded onto a workstation (Advantage Workstation VolumeShare 4 XT, GE Healthcare, Milwaukee, WI) and converted into three-dimensional (3D) images (Fig. 1) using the workstation's onboard software.

Two radiologists classified the cases based on data from the 3D images, and those with poorly visible lymphatic routes or lymph nodes were also classified using the original CT image as a reference.

### **Anatomical classification procedure**

SLNs were first classified into four categories based on the report of Yamamoto et al (Yamamoto et al. 2016) and then each category was subdivided as follows: single route/single SLN was subdivided into (a) no branching, (b) breast side branching, (c) axillary side branching, (d) breast and axillary sides branching, and (e) other; single route/multiple SLNs into (a) breast side branching, (b) axillary side branching, and (c) breast and axillary sides branching; multiple routes/single SLN into (a) no branching in each route, (b) mixed, with and without branching, and (c) each route with separate branching; and multiple

routes/multiple SLNs into (a) no branching, (b) branching on each route, and (c) branching and flow from other route.

Based on the reports of Kutsuna and Kaneko (Kutsuna 1968; Kaneko 2001), SLNs were classified into three categories, based on the flow direction: (a) flowing outward on the superficial fascia toward the axillary lymph nodes, (b) flowing upward on the superficial fascia toward the infraclavicular or supraclavicular nodes, and (c) flowing toward the midline and reaching the parasternal or contralateral axillary lymph nodes.

Regional lymph nodes involved in breast cancer include the axillary, brachial, subpectoral, and infraclavicular lymph nodes. The axillary lymph nodes were further classified into (a) upper pectoral lymph nodes, (b) central axillary lymph nodes, and (c) lower pectoral lymph nodes. They were classified based on their location compared with rib level into (a) the upper pectoral lymph nodes located between the upper border of the second rib and the lower border of the third rib, (b) central axillary lymph nodes located above (a), and (c) the lower pectoral lymph nodes located between the fourth and sixth ribs.

### **Statistical analysis**

Possible correlations between anatomical classification (good vs. poor imaging) and histological metastasis (presence vs. absence) were tested using the Chi-squared test with a 5% significance level. The test was performed using Excel 2013 software (Microsoft Corporation, Redmond, WA) with the Statcel 4

add-in package.

## Results

Patients' characteristics are listed in Table 1. Among the 464 patients, anatomical classification could be performed in 434 (93.5%), but not in 30 (6.5%). Although these 30 cases were impossible to classify, there were seven cases where only the lymphatic routes were rendered, four where only the SLNs were drawn, and 19 were without rendering. Of the 434 cases that permitted anatomical classification, 99 (22.8%) showed histological evidence of metastases, whereas 335 (77.2%) showed no metastases. Of the 30 cases that could not be anatomically classified, four (13.3%) showed histological evidence of metastases, whereas 26 (86.7%) showed no metastases. There was no correlation between anatomical classification (good vs. poor imaging) and metastases (presence vs. absence) ( $p = 0.22$ ).

Among the 434 cases analyzed, the largest number of cases showed single route/single SLN ( $n = 296$ , 68.2%), followed by multiple routes/multiple SLNs ( $n = 59$ , 13.6%), single route/multiple SLNs ( $n = 53$ , 12.2%), and multiple routes/single SLN ( $n = 26$ , 6.0%) (Fig. 2). In the 296 cases with single route/single SLN, the largest number of cases showed no branching ( $n = 145$ , 49.0%) (Fig. 3). Of the 53 cases with single route/multiple SLNs, most showed axillary side branching ( $n = 31$ , 58.5%) (Fig. 4). Of the 26 cases with multiple routes/single SLN, 12 (46.2%) showed no branching in each route, 12 (46.2%) showed mixed, with and without branching, and 2 (7.7%) showed each route with separate branching (Fig. 5). Of the 59 cases with multiple routes/multiple SLNs, 26 (44.1%) showed no branching and comprised the largest group (Fig. 6), whereas four (6.8%) had routes that did not reach the axilla. Furthermore,

branching on each route was seen in 23 cases (39.0%), and among these, 14 (23.7%) showed branching on either route, and nine (15.3%) showed branching into multiple SLNs.

The 434 cases were also categorized according to the flow direction of SLNs, wherein 429 (98.8%) were found to be flowing outward on the superficial fascia toward axillary lymph nodes, three (0.7%) toward the midline that reached the parasternal lymph nodes, and zero flowing upward on the superficial fascia toward the infraclavicular or supraclavicular lymph nodes (Fig. 7). Additionally, there was one case each (0.2%) of flow that reached the interpectoral and supraclavicular lymph nodes via the axilla.

Next, all 434 cases were classified according to the rib height at which the SLNs were present; 323 (74.4%) were upper pectoral lymph nodes, 54 (12.4%) were central axillary lymph nodes, and 22 (5.1%) were lower pectoral lymph nodes. In addition, there were 25 cases (5.8%) where the central axillary lymph nodes were merged with the upper pectoral lymph nodes, nine (2.1%) where the upper and lower pectoral lymph nodes were merged, and one case (0.2%) with both of the above-mentioned phenomena (Fig. 8).

## Discussion

The current standard of breast cancer surgery recommends ALND only if the results of SLN biopsy warrant it (Krag et al. 2010; The Japanese Breast Cancer Society 2016; Lyman et al. 2005; Lyman et al. 2014). Furthermore, ALND may be completely omitted because the prognosis remains unchanged despite the SLNs being positive for micrometastases ( $< 2$  mm but  $> 0.2$  mm) (Galimberti et al. 2013; Maaskant-Braat et al. 2011). Some studies have explored this option in macrometastases ( $> 2$  mm) (Giuliano et al. 2011; Agresti et al. 2014; Donker et al. 2014), however, it is still uncertain if ALND is completely unnecessary. Thus, it is important to completely understand the anatomy and number of lymphatic routes and SLNs and their location.

A total of 480 cases that were clinically diagnosed with node-negative breast cancer underwent preoperative SLN CT-LG. Of these, 16 (6, previous ipsilateral breast surgery; 10, preoperative chemotherapy) were excluded from this study because Lizarraga et al. (Lizarraga et al. 2013) reported that SLNs were poorly visualized after surgery on the ipsilateral chest or axilla. Van der Ploeg et al. (Van der Ploeg et al. 2010) reported that compared with untreated breasts, lymphatic flow in a previously treated breast occurs less often to the axilla and more often to nodes elsewhere, as far away as the contralateral side. In the present study, there was no lymphatic flow to the extra axilla, but five of the six surgery cases showed poor visualization. Previous reports also suggested that chemotherapy interferes with subsequent scintigraphy and lowers the SLN detection rate (Kuehn et al. 2013). Accordingly, 2 of the

10 cases that underwent preoperative chemotherapy showed poor visualization.

Thirty cases (6.5%) were excluded from anatomical classification due to poor imaging, and this proportion was similar to those of previous reports (Yamamoto et al. 2016; Takahashi et al. 2008). The causes of poor imaging can be classified into two categories: (1) the contrast agent does not flow into the lymphatic routes and lymph nodes, resulting in no capture; and (2) the contrast agent flows out from the lymphatic routes or lymph nodes before the scan. Probable reasons for the obstruction of contrast agent include flow occlusion by tumor cells in the lymphatic routes and lymph nodes (Lehman et al. 2013). However, using the Chi-squared test, there was no correlation between image quality and metastasis ( $p = 0.22$ ). The flowing out of the contrast agent could be attributed to the agent being washed out before the scan, possibly due to increases in the number and diameter of lymphatic routes and the lymphatic network formed due to tumor-mediated lymphangiogenesis (Ran et al. 2010). Another possible reason for the flowing out of the contrast agent is the use of a water-soluble agent instead of an oil contrast one, such as ethyl ester of iodinated poppy seed oil fatty acid (Gómez et al. 2012). Although poor visualization can result from problems in the procedure, it is difficult to believe that these two are substantially related, because the injection of the contrast agent and scanning are sequentially and immediately performed. However, the reasons for such poor imaging should be investigated furthermore.

Several injection techniques for SLN detection have been used to date, such as peritumoral or intranipple injection. In previous studies (Suga et al. 2003; Minohata et al. 2011), injection directly into the nipple was performed because it allows easy observation of the morphological and anatomical



features, such as the number of lymphatic routes and lymph nodes, and it avoids the use of image-guided injections of nonpalpable breast lesions. On the other hand, Suami et al. (Suami et al. 2009b) stated that compared with peritumoral injections, radioisotope intranipple injections alone may not detect some SLNs because numerous valves regulate the lymphatic flow and prevent its backflow. However, their findings are not necessarily applicable to the present study because of the variations in assessing dead and living bodies. Blumgart et al. (Blumgart et al. 2011a) reported that lymphatic flow should be evaluated without peritumoral injections because peritumoral injections of radioisotope can cause a large region of breast tissue to contain radioactive tracer, thereby making observations less precise. Several other reports have stated that SLN identification is not affected by the injection site regardless of whether the injection is intranipple or peritumoral (Kern 2001; Caruso et al. 2014; Rosenow et al. 2012). Therefore, the present study used intranipple injection.

From the results of this study, single route/single SLN was the predominant category, and single route/multiple SLNs and multiple routes/multiple SLNs categories comprised 10% of the cases; these results were similar to those reported by Yamamoto et al (Yamamoto et al. 2016). The prevalence of multiple routes/single SLN was 6.0% in the present study and 11% in the report of Yamamoto et al, however, there was only a marginal difference between them.

In the present study, Yamamoto's four categories of breast CT-LG were further subdivided based on branching patterns and the height at which the branching occurred. Such subdivision showed that lymphatic networks were complicated, and that CT-LG was useful in detecting such anatomical

complexity.

Kutsuna and Kaneko (Kutsuna 1968; Kaneko 2001) reported cases where the lymph flowed into the parasternal lymph node, into the infraclavicular or supraclavicular lymph nodes without going through the axilla, and into the axilla of the contralateral side. There was no case of lymph flow into the infraclavicular or supraclavicular lymph nodes without going through the axilla, and into the axilla of the contralateral side in the present study. The lymphatic flow to the parasternal lymph node and to the supraclavicular lymph node via the axilla were observed.

According to the anatomical definition, the axilla refers to the indentation under the shoulder joint (Moore et al. 2014). The region of the axillary lymph nodes in breast cancer is wide and extends not only to the axillary locus, but also to the lateral chest and surrounding clavicle. In agreement with this fact, it was clearly shown that the axillary SLNs were widely distributed, and that a focus restricted only to the axillary locus could lead to potential oversight. To avoid such omissions, it is important to understand the location of SLNs. Therefore, CT-LG was used for easy classification of SLN location. While some SLNs can be identified using blood vessels and muscles as landmarks, the central axillary and pectoral lymph nodes, especially those distributed from the axillary region to the lateral chest, are difficult to locate. However, the ribs can be used to facilitate the location of these lymph nodes; pectoral lymph nodes can be divided into the upper group (between the second and third ribs) and the lower group (between the fourth and sixth ribs) (Akiyama et al. 2015), whereas the central axillary lymph nodes are distributed above the upper group. Furthermore, since CT-LG was performed in the same posture as that during surgery, SLN

location should not vary, which is considered very useful.

In conclusion, a wide variation in the number of lymphatic routes and their branching patterns, number of SLNs and their locations, and the flow direction of SLNs was seen using CT-LG. CT-LG is useful for detecting lymphatic routes and SLNs in breast cancers and understanding SLN morphology, and it can be easily implemented in settings with ordinary CT. A better understanding of the anatomical morphology of lymphatic routes and SLNs preoperatively is considered clinically very important and useful for the treatment of breast cancers.

## **Acknowledgments**

The authors thank Fumiyasu Tsushima, Shinya Kakehata, Hiromasa Fujita, Mitsuki Tanaka, Saya Kakizaki, Soichiro Tatsuo, Sayuri Tatsuo, and Akiko Igawa for collecting the amount number of data.

## **Compliance with ethical standards**

**Conflict of Interest:** The authors declare that they have no conflict of interest.

**Ethical statement:** This retrospective study was approved by our ethics committee, and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

**Informed consent:** Informed consent for using case information was obtained from all patients.

## References

- Agresti R, Martelli G, Sandri M et al (2014) Axillary lymph node dissection versus no dissection in patients with T1N0 breast cancer: a randomized clinical trial (INT09/98). *Cancer* 120:885–893. doi: 10.1002/cncr.28499
- Akiyama K, Imai S, Ogawa S et al (2015) Keibu, Ekika, Kyō-heki rinpa-seru (Cervical, axillary and thoracic lymph nodes). In: Kumita S (ed) *Gazō-sindan no tameno Color Atlas* (Color Atlas for diagnostic imaging). Kanehara shuppan, Tokyo, pp 10–11
- Blumgart EI, Uren RF, Nielsen PMF et al (2011a) Lymphatic drainage and tumor prevalence in the breast: a statistical analysis of symmetry, gender and node field independence. *J Anat* 218:652–659. doi: 10.1111/j.1469-7580.2011.03162.x
- Blumgart EI, Uren RF, Nielsen PMF et al (2011b) Predicting lymphatic drainage patterns and primary tumour location in patients with breast cancer. *Breast Cancer Res Treat* 130:699–705. doi: 10.1007/s10549-011-173-2
- Caruso G, Cipolla C, Costa R et al (2014) Lymphoscintigraphy with peritumoral injection versus lymphoscintigraphy with subdermal periareolar injection of technetium-labeled human albumin to identify sentinel lymph nodes in breast cancer patients. *Acta Radiol* 55:39–44. doi: 10.1177/0284185113493775
- Donker M, van Tienhoven G, Straver ME et al (2014) Radiotherapy or surgery of the axilla after a

positive sentinel node in breast cancer (EORTC 10981-22023 AMAROS): a randomized, multicentre, open-label, phase 3 non-inferiority trial. *Lancet Oncol* 15:1303–1310. doi:

10.1016/S1470-2045(14)70460-7

Galimberti V, Cole BF, Zurrada S et al (2013) Axillary dissection versus no axillary dissection in patients with sentinel-node micrometastases (IBCSG 23-01): a phase 3 randomized controlled trial. *Lancet Oncol* 14:297–305. doi: 10.1016/S1470-2045(13)70035-4

Giuliano AE, Hunt KK, Ballman KV et al (2011) Axillary dissection vs no axillary dissection in women with invasive breast cancer and sentinel node metastasis: a randomized clinical trial. *JAMA* 305:569–575. doi: 10.1001/jama.2011.90.

Gómez FM, Martínez-Rodrigo J, Martí-Bonmatí L et al (2012) Transnodal lymphangiography in the diagnosis and treatment of genital lymphedema. *Cardiovasc Intervent Radiol* 35:1488–1491. doi: 10.1007/s00270-012-0445-6

Kaneko U (2001) Rinpa-kei (Lymphatic system). In: Kaneko U (ed) *Nihon jintai kaibō-gaku ge-kan* (Japanese anatomy of the second volume). Nanzando, Tokyo, pp 198–207

Kern KA (2001) Lymphoscintigraphic anatomy of sentinel lymphatic channels after subareolar injection of technetium 99m sulfur colloid. *J Am Coll Surg* 193:601–608. doi: org/10.1016/S1072-7515(01)01068-7

Krag DN, Anderson SJ, Julian TB et al (2010) Sentinel-lymph-node resection compared with conventional axillary-lymph-node dissection in clinically node-negative patients with breast cancer:

overall survival findings from the NSABP B-32 randomized phase 3 trial. *Lancet Oncol* 11:927–933. doi:

10.1016/S1470-2045(10)70207-2

Kuehn T, Bauerfeind I, Fehm T et al (2013) Sentinel-lymph-node biopsy in patients with breast cancer

before and after neoadjuvant chemotherapy (SENTINA): a prospective, multicenter cohort study. *Lancet*

*Oncol* 14:609–618. doi: 10.1016/S1470-2045(13)70166-9

Kutsuna M(1968) Jōsi oyobi kyōheki no rinpasetu (Lymph nodes of upper limbs and chest walls). In:

Kutsuna M (ed) *Nihon-jin no rinpakei-kaibō-gaku* (Japanese lymphatic anatomy). Kanehara shuppan,

Tokyo, pp 113–114

Lehman HL, Dashner EJ, Lucey M et al (2013) Modeling and characterization of inflammatory breast

cancer emboli grown in vitro. *Int J Cancer* 132:2283–2294. doi: 10.1002/ijc.27928

Lizarraga IM, Scott-Conner CEH, Muzahir S, Weigel RJ, Graham MM, Sugg SL (2013) Management of

contralateral axillary sentinel lymph nodes detected on lymphoscintigraphy for breast cancer. *Ann Surg*

*Oncol* 20:3317–3322. doi: 10.1245/s10434-013-3151-y

Lyman GH, Giuliano AE, Somerfield MR et al (2005) American society of clinical oncology guideline

recommendations for sentinel lymph node biopsy in early-stage breast cancer. *J Clin Oncol* 23:7703–7720.

doi: 10.1200/JCO.2005.08.001

Lyman GH, Temin S, Edge SB et al (2014) Sentinel lymph node biopsy for patients with early-stage

breast cancer: American society of clinical oncology clinical practice guideline update. *J Clin Oncol*

32:1365–1383. doi: 10.1200/JCO.2013.54.1177

Maaskant-Braat AJ, van de Poll-Franse LV, Voogd AC et al (2011) Sentinel node micrometastases in breast cancer do not affect prognosis: a population-based study. *Breast Cancer Res Treat* 127:195–203. doi: 10.1007/s10549-010-1086-6

Minohata J, Takao S, Hirokaga K (2011) Sentinel lymph node biopsy using CT lymphography in breast cancer. *Breast Cancer* 18:129–136. doi: 10.1007/s12282-009-0183-z

Moore KL, Dalley AF, Agur AMR (2014) Axilla. In: Taylor C (ed) *Clinically oriented anatomy* seventh edition. Wolters Kluwer/Lippincott Williams and Wilkins, Philadelphia, p 173

Ran S, Volk L, Hall K, Flister MJ (2010) Lymphangiogenesis and lymphatic metastasis in breast cancer. *Pathophysiology* 17:229–251. doi: 10.1016/j.pathophys.2009.11.003

Rosenow T, Biedendieck C, Fricke H et al (2012) Sentinel lymph node biopsy in patients with breast cancer: comparison of peritumoral and periareolar injection. *Geburtshilfe Frauenheilkd* 72:1024–1028. doi: 10.1055/s-0032-1315304

Suami H, Pan W-R, Taylor GI (2009a) Historical review of breast lymphatic studies. *Clin Anat* 22:531–536. doi: 10.1002/ca.20812

Suami H, Pan W-R, Taylor GI (2009b) Redefined lymphatic anatomy of the breast with clinical implications. In: Leong SPL (ed) *Current Clinical Oncology*. Humana Press, Berlin, pp 45-55. doi: 10.1007/978-1-60327-087-8\_5

Suga K, Ogasawara N, Okada M, Matsunaga N (2003) Interstitial CT lymphography-guided localization of breast sentinel lymph node: preliminary results. *Surgery* 133:170–179. doi: 10.1067/msy.2003.17



Takahashi M, Sasa M, Hirose C et al (2008) Clinical efficacy and problems with CT lymphography in identifying the sentinel node in breast cancer. *World J Surg Oncol*. doi: 10.1186/1477-7819-6-57

The Japanese breast cancer society (2016) The Japanese breast cancer society clinical practice guideline for surgical treatment of breast cancer, 2015 edition. *Breast Cancer* 23:367–377. doi:

10.1007/s12282-016-0671-x

Uren RF, Howman-Giles R, Thompson JF (2003) Patterns of lymphatic drainage from the skin in patients with melanoma. *J Nucl Med* 44:570–582

Van der Ploeg IMC, Oldenburg HSA, Rutgers EJT et al (2010) Lymphatic drainage patterns from the treated breast. *Ann Surg Oncol* 17:1069-1075. doi: 10.1245/s10434-009-0841-6

Yamamoto S, Suga K, Maeda K, Maeda N, Yoshimura K, Oka M (2016) Breast sentinel lymph node navigation with three-dimensional computed tomography-lymphography: a 12-year study. *Breast Cancer* 23:456–462. doi: 10.1007/s12282-015-0584-0

## Figure legends

**Fig. 1** A 3D-CT lymphogram of a woman in her 50s with left breast cancer

A single lymphatic route (thin arrow) from the periareolar area (thick arrow) drains into a single SLN (arrowhead).

**Fig. 2** Anatomical classification of SLNs in 434 cases drawing with reference to the reports of Yamamoto et al. (Yamamoto et al. 2016).

**a** Single route/single SLN (n = 296, 68.2%). **b** Single route/multiple SLNs (n = 53, 12.2%). **c** Multiple routes/single SLN (n = 26, 6.0%). **d** Multiple routes/multiple SLNs (n = 59, 13.6%).

**Fig. 3** Further classification results of single route/single SLN in 296 cases

**a** No branching (n = 145, 49.0%). **b** Breast side branching (n = 84, 28.4%). **c** Axillary side branching (n = 35, 11.8%). **d** Breast and axillary sides branching (n = 25, 8.4%). **e** Other (n = 7, 2.4%).

**Fig. 4** Further classification results of single route/multiple SLNs in 53 cases

**a** Breast side branching (n = 3, 5.7%). **b** Axillary side branching (n = 31, 58.5%). **c** Breast and axillary sides branching (n = 19, 35.8%).

**Fig. 5** Further classification results of multiple routes/single SLN in 26 cases

**a** No branching in each route (n = 12, 46.2%). **b** Mixed, with and without branching (n = 12, 46.2%). **c** Each route with separate branching (n = 2, 7.7%).

**Fig. 6** Further classification results of multiple routes/multiple SLNs in 59 cases

**a** No branching (n = 26, 44.1%). **b** Branching on each route (n = 23, 39.0%). **c** Branching and flow from

other route (n = 10, 16.9%).

**Fig. 7** Classification results by SLN flow direction

**a** Flowing outward on the superficial fascia toward the axillary lymph nodes (n = 429, 98.8%). **b** Flowing upward on the superficial fascia toward the infraclavicular or supraclavicular nodes (n = 0, 0%). **c** Flowing toward the midline and reaching the parasternal lymph nodes (n = 3, 0.7%). **d** Flow that reached the interpectoral lymph node via the axilla (n = 1, 0.2%). **e** Flow that reached the supraclavicular lymph node via the axilla (n = 1, 0.2%).

**Fig. 8** Classification results based on SLN location compared with rib level

**a** Upper pectoral lymph nodes (n = 323, 74.4%). **b** Central axillary lymph nodes (n = 54, 12.4%). **c** Lower pectoral lymph nodes (n = 22, 5.1%). **d** Merged central axillary lymph nodes and upper pectoral lymph nodes (n = 25, 5.8%). **e** Merged upper and lower pectoral lymph nodes (n = 9, 2.1%). **f** Both of the above-mentioned phenomena (n = 1, 0.2%).

**Table 1.** Patients’ background characteristics

Parameter	Values n (%)
Number of patients	464
Age in years, average (range)	56 (24-88)
Sex	
Male	0
Female	464 (100%)
SLN metastasis	
Positive	104 (22.4%)
Negative	360 (77.6%)
Histological type	
DCIS	62 (13.4%)
Papillotubular ca.	110 (23.7%)
Solid-tubular ca.	50 (10.8%)
Scirrhouc ca.	208 (44.8%)
Mucinous ca.	12 (2.6%)
Invasive micropapillary ca.	7 (1.5%)

---

Invasive lobular ca.	6 (1.3%)
Apocrine ca.	6 (1.3%)
Medullary ca.	3 (0.6%)

---

*DCIS* ductal carcinoma in situ, *Ca.* carcinoma

Fig. 1




Fig. 2




Fig. 3









Fig. 4



Breast side  
branching  
3 cases  
(5.7%)




Axillary side  
branching  
31 cases  
(58.5%)



Breast and  
axillary sides  
branching  
19 cases  
(35.8%)

Fig. 5



No branching  
in each route  
12 cases  
(46.2%)




Mixed, with and  
without branching  
12 cases  
(46.2%)




Each route with  
separate branching  
2 cases  
(7.7%)


Fig. 6



No branching  
26 cases  
(44.1%)



Branching on  
each route  
23 cases  
(39.0%)



Branching and  
flow from other route  
10 cases  
(16.9%)

Fig. 7




Fig. 8

